


Ebola Mandatory Protocol

Algorithm for OB Patient Screening


Oct. 22, 2014 5 p.m.

Patient Arrives at Labor & Delivery



Revised: 10/22/2014 5 p.m. 1 of 2

Patient Arrives at Labor & Delivery (continued from page 1)



Do not touch the patient until **Treatment PPE** is donned per protocol

- Once Treatment PPE is donned, perform vaginal delivery and keep infant newborn with mom for normal newborn care, or place in second room for NICU treatment (Do not take infant to newborn or NICU)
- If Cesearian section is necessary:
 - -Don Treatment PPE
- -Place mask on patient
- -Clear path to ICU, per Intra-transport process
- OB performs Cesearian section utilizing least number of personnel to safely care for patient and infant (ex. L&D RN, ICU RN)
- Have NICU RN or Neonatal Nurse Practitioner donned in Treatment PPE outside of room in case intensive intervention with infant becomes necessary
- -Patient stays in ICU room and infant stays with mom unless intensive care is needed for the infant
- -There should be an ISTAT for each patient
- -If infant needs NICU care, place in second room (Do not take infant to newborn or NICU)
- Notify regarding need to decontaminate hallway and ICU. Secure room until decontamination is complete.

Revised: 10/22/2014 5 p.m. 2 of 2